



Dehydrated Culture Media
Bases / Media Supplements

Technical Information

Antifungal Assay Agar

Product Code: DM 1164

Application: Antifungal Assay Agar is recommended for assaying antifungal activity of pharmaceutical products and other materials by the cylinder plate or disc diffusion method.

Composition**

Ingredients	Gms / Litre
Dextrose	50.000
Sodium citrate	4.500
Potassium phosphate	0.550
Citric acid	1.000
Casein enzymic hydrolysate	4.000
Pyridoxine hydrochloride	0.00025
Thiamine	0.00025
Inositol	0.025
Calcium pantothenate	0.0025
Niacin	0.0025
Potassium chloride	0.425
Calcium chloride	0.125
Magnesium sulphate	0.125
Ferric chloride	0.0025
Manganese sulphate	0.0025
Biotin	0.000008
Agar	15.000
Final pH (at 25°C)	5.5±0.2

**Formula adjusted, standardized to suit performance parameters

Principle & Interpretation



Fungal infections have been increased dramatically in the past decade, as systemic infections or as co-infections in the immune compromised patients & individual suffering from cancer or AIDS ^(1,2). Unfortunately, due to the limited number of antifungal drugs currently available, fungal infections tend to develop resistance to these drugs very rapidly. For these reasons, fungal infections now days are responsible for much higher mortality rates than bacterial infections ⁽³⁾. This led to the need for development of rapid and accurate methods for antifungal screening and susceptibility testing. Antifungal Assay Agar was formulated by Berger and Lazecka for assaying antifungal activity of pharmaceutical products and other materials by both base and seed layer assays using cylinder plate or disc diffusion methods. The defined ingredients in the medium provide the necessary nutrients and growth factors required for the development of the test culture. Phosphate is included in this medium for good buffering action. Dextrose in the medium serves as a carbon and energy source. Other ingredients like the sulphates; vitamins, growth factors etc are added to enhance the growth of the test organisms, so that the inhibition obtained is always due to the antifungal agents and not due to nutrient depletion. **Cylinder plate method:** This method was first devised by Abraham et al ⁽⁴⁾ which was later modified by Schmidt and Moyer ⁽⁵⁾ and It depends upon diffusion of the antibiotic from vertical steel cylinders placed on the surface of inoculated agar medium. This produces zones of inhibition around the cylinder containing antibiotic solution depending upon the concentration of the antibiotic in the cylinder. This method is commonly employed in the assay of pharmaceutical preparations of Penicillin and

other antibiotics. For assay, Petri plates with 20 x100 mm dimension and stainless steel or porcelain cylinders with the outside diameter 8 mm, inside diameter 6 mm and length 10 mm and used. All dimensions should have a tolerance of 0.1 mm. The cylinders should be carefully cleaned to remove all the impurities. For assays requiring base and seed layer, the base layer is allowed to solidify first and then overlaid with the seed agar containing the proper concentration of the test organism. Most assays require base layer of 21 ml and seed layer of 4 ml. Generally 6 cylinders are used per plate. The cylinders are placed on inoculated plates at equal distance.

Paper-disc method: Paper discs with a diameter of 9 mm are impregnated with the antibiotic solution and placed on the culture medium. Antibiotic can also be applied to the disc after it has been placed on the medium. Plates containing a single layer of medium with 2 mm thickness may be used for these tests. All other steps are similar to the cylinder plate method.

Methodology

Suspend 75.76 grams of powder media in 1000 ml distilled water. Shake well and heat to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Mix well and pour into sterile Petri plates.

Quality Control

Physical Appearance

Cream to beige homogeneous free flowing powder

Gelling

Firm, comparable with 1.5% Agar gel

Colour and Clarity of prepared medium

Light yellow coloured clear to slightly opalescent gel forms in Petri plates

Reaction

Reaction of 7.57% w/v aqueous solution at 25°C. pH : 5.5±0.2

pH Range:-

5.30-5.70

Cultural Response/Characteristics

DM1164: Cultural characteristics observed after an incubation at 25-30°C for 18-48 hours.

Organism	Inoculum (CFU)	Growth	Recovery
<i>Saccharomyces cerevisiae</i> ATCC 9763	50-100	luxuriant	>=70%



Dehydrated Culture Media
Bases / Media Supplements

**Aspergillus brasiliensis* 50-100 Luxuriant
ATCC 16404
Key * - Formerly known as *Aspergillus niger*

Storage and Shelf Life

Dried Media: Store below 30°C in tightly closed container and use before expiry date as mentioned on the label.

Prepared Media: 2-8° in sealable plastic bags for 2-5 days.

Further Reading

1. Beck-Sague C. and Jarvis W. R., 1993, J. Infect. Dis., 167:1247-1251.
2. Berrouane Y. F., Herwaldt L. A., and Pfaller M. A., 1999, J. Clin. Microbiol., 37:531-537.
- Weinstein, M. P., Towns M. L., Quartey S. M., Mirrett S., Reimer L. G., Parmigiani G. and Reller L. B., 1997., Clin. Infect. Dis., 24:584-602.
4. Abraham, Chain, Fletcher, Florey, Gardner, Heatley and Jennings, 1941, Lancet ii: 177.
5. Schmidt and Moyer, 1944, J. Bacteriol., 47:199.

Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conform solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at **CDH** is true and accurate
- **Central Drug House Pvt. Ltd.** reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents. Do not use the products if it fails to meet specifications for identity and performance parameters.

